www.mikedemy.com

3.1 Reciprocal Trigonometric Functions

Question Paper

| Course | CIEASMaths |
|------------|--|
| Section | 3. Trigonometry |
| Торіс | 3.1 Reciprocal Trigonometric Functions |
| Difficulty | Hard |

| Time allowed: | 40 |
|---------------|------|
| Score: | /28 |
| Percentage: | /100 |

Question la

(a) Rewrite $\tan \theta \csc \theta$ as a single trigonometric function.

[2 marks]

Question 1b

(b) Hence solve, in the range $-\pi < \theta \leq \pi$, the equation

$$\tan\theta\csc\theta = -\frac{2\sqrt{3}}{3}.$$

[3 marks]

Question 2

Solve, in the range $0 \le \theta \le 2\pi$, the equation

$$\frac{2}{\csc \theta} - \csc \theta = 1.$$

[6 marks]

Question 3

Using the double angle formula $\sin 2A \equiv 2 \sin A \cos A$, find the solutions to the equation

$$\sec x \csc x - 75 = 5 \csc 2x$$

in the range $-\pi < x \le \pi$. Give your answers correct to 3 significant figures.

[6 marks]

Question 4a

(a) Show that the equation

 $2 \cot^2 x = 1 - 5 \operatorname{cosec} x$

can be rewritten in the form

$$(2\csc x - 1)(\csc x + 3) = 0.$$

[3 marks]

Question 4b

(b) Hence solve, in the range $0 \le x \le 2\pi$, the equation

 $2 \cot^2 x = 1 - 5 \operatorname{cosec} x$

giving your answers correct to 3 significant figures.

[3 marks]

Question 5

- (i) Sketch, in the interval $-2\pi \le \theta \le 2\pi$, the graph of $y = -5 + \frac{1}{2} \sec \theta$, include asymptotes and label the coordinates of all maximum and minimum points.
- (ii) Hence deduce the range of values for k for which the equation $-5 + \frac{1}{2} \sec \theta = k$ has no solutions.

[5 marks]